

CENTRAL INTELLIGENCE AGENCY
INFORMATION REPORT

REPORT

CD NO.

COUNTRY East Germany

DATE DISTR. 9 March 1954

SUBJECT Development of Cadmium Sulfide Crystals at
the Institute for Medicine and Biology, Berlin-Buch

NO. OF PAGES 1

016043

PLACE
ACQUIREDNO. OF ENCLS.
(LISTED BELOW)

25X1

DATE OF
INFO.SUPPLEMENT TO
REPORT NO.THIS DOCUMENT CONTAINS INFORMATION AFFECTING THE NATIONAL DEFENSE
OF THE UNITED STATES, WITHIN THE MEANING OF TITLE 18, SECTIONS 793
AND 794, OF THE U. S. CODE, AS AMENDED. ITS TRANSMISSION OR REVEL-
ATION OF ITS CONTENTS TO OR RECEIPT BY AN UNAUTHORIZED PERSON
IS PROHIBITED BY LAW. THE REPRODUCTION OF THIS FORM IS PROHIBITED.

THIS IS UNEVALUATED INFORMATION

25X1

25X1

1. The ~~Department~~ of the Academy Institute for Medicine and Biology in Berlin-Buch is working on the development of cadmium sulfide crystals for X-ray dosimetry. The work is being carried out by Physicist Krumbiegel (fnu) in the laboratory of Dr. Liselott Herforth. The crystals are produced by vaporizing cadmium in hydrogen sulfide at a temperature of 1,000° Centigrade.

2. Two methods of crystal activation are used:

- a. Fine silver is vaporized at the same time that the cadmium is exposed to the hydrogen sulfide stream. The silver diffuses into the crystals and ultimately improves their conductivity.
- b. The finished crystals are tempered at a temperature between 500 and 600° Centigrade with addition of fine silver.

Experiments carried out thus far seem to indicate that method b. furnishes better results than method a. It is expected, however, that the experiments will have to be continued for a period of six more months before an ultimate decision on the preferability of either of the two methods can be reached.

3. Before the crystal is exposed to an X-ray source it is inserted into a circuit so that a voltage between 10 and 40 Volts is laid on the crystal. Higher voltages make the crystal unworkable. The voltage has to be gradually increased from 10 to a maximum of 40 Volts. The current must not exceed 200 micro-amperes because higher currents also make the crystal unuseable. The initial X-ray doses to which the crystal is exposed must be weak; later doses are gradually increased.

25 YEAR RE-REVIEW

25X1

CLASSIFICATION: SECRET

STATE	<input checked="" type="checkbox"/>	NAVY	<input checked="" type="checkbox"/>	NSRB	<input checked="" type="checkbox"/>	DISTRIBUTION	<input checked="" type="checkbox"/>	OSI Ev	<input checked="" type="checkbox"/>
ARMY	<input checked="" type="checkbox"/>	ALL	<input checked="" type="checkbox"/>						